
GENERATION OF PHOSPHOROUS ESTER PHOSPHORUS ANIDE AND PHOSPHINE DERIVATIVE. (U) DUKE UNIV DURHAM NC DEPT OF CHEMISTRY LD QUIN 17 OCT 86 ARO-22945.7-CH AD-A175 747 1/1 UNCLASSIFIED NL





		~				
	REPORT DOCUM	MENTATION	PAGE			
AD_ A175 747	1b. RESTRICTIVE MARKINGS					
AD-A175 747		3. DISTRIBUTION/AVAILABILITY OF REPORT				
B. DECLASSIFICATION / WITHING COMMISSION	Approved	i for public	c release	;		
	distribution unlimited.					
I. PERFORMING ORGANIZATION REPORT NUMBE	R(S)	5. MONITORING	ORGANIZATION R	EPORT NUMB	ER(S)	
			22945.7-			
60. NAME OF PERFORMING ORGANIZATION 6b. OFFICE SYMBOL (If applicable)		7a. NAME OF MO	ONITORING ORGA	NIZATION		
Duke University		U. S. A	rmy Researc	h Office		
Sc. ADDRESS (City, State, and ZIP Code)		7b. ADDRESS (Cit	-	Code)		
Department of Chemistry, Paul Chemical Lab, Durham, NC 27			ox 12211 h Triangle	Park NC	27700_22	
		<u>]</u>				
Bo. NAME OF FUNDING/SPONSORING ORGANIZATION	8b. OFFICE SYMBOL (If applicable)	9. PROCUREMENT	T INSTRUMENT ID	ENTIFICATION	NUMBER	
U. S. Army Research Office	<u> </u>		AL-3-86-K-00			
Bc. ADDRESS (City, State, and ZIP Code)		10. SOURCE OF F	PROJECT	rs Task	WORK U	
P. O. Box 12211 Research Triangle Park, NC 23	7709-2211	PROGRAM ELEMENT NO.	NO.	NO.	ACCESSIO	
Coordination Number 12 PERSONAL AUTHOR(S) Louis D. Qu 13a TYPE OF REPORT 13b TIME CO		14 DATE OF REDO	ORI (Year Month	Dav) hs p	AGE COUNT	
Final From 12/	OVERED /10/85 TO 7/31/86	1986 (October 17	, Jey) 13. P/	AGE COUNT 5 pages	
of the author(s) and should not bolicy or decision unless so 17 COSATI CODES FIELD GROUP SUB-GROUP	opinions and/or t be construed a designated by 18. SUBJECT TERMS	s an officia	l Departmen	nt of the	Army posi	
	and identify by black	number)		-		
19 ABSTRACT (Continue on reverse if necessary	and identity by block		systems have	e been pr	epared as	
Derivatives of the 5-pho	sphabicyclo[2.2	.2]octene-2 : . Metaphosph	nate generat	cors are a	also heino	
Derivatives of the 5-pho precursors on thermolysis of prepared, especially with lar	sphabicyclo[2.2 phosphalkenenes	. Metaphospl	nate generat provide stat	cors are a pilization	also being n of the	
Derivatives of the 5-pho precursors on thermolysis of	sphabicyclo[2.2 phosphalkenenes	. Metaphospl	nate generat provide stab	oilization	n of the	
Derivatives of the 5-pho precursors on thermolysis of prepared, especially with lar	sphabicyclo[2.2 phosphalkenenes	. Metaphospl	nate generat provide stab	oilization	n of the	
Derivatives of the 5-pho precursors on thermolysis of prepared, especially with lar	sphabicyclo[2.2 phosphalkenenes	. Metaphospl	nate generat provide stab	cors are a	n of the	
Derivatives of the 5-pho precursors on thermolysis of prepared, especially with lar	sphabicyclo[2.2 phosphalkenenes ge bulky groups	. Metaphospl	nate generat	cors are a	n of the	
Derivatives of the 5-pho precursors on thermolysis of prepared, especially with lar species. DTIC FILE C	osphabicyclo[2.2 phosphalkenenes ge bulky groups	Metaphosph that might p	provide stat	oilization	n of the	
Derivatives of the 5-pho precursors on thermolysis of prepared, especially with lar species.	osphabicyclo[2.2 phosphalkenenes ge bulky groups	Metaphosph that might p	ECURITY CLASSIFI nclassified (include Area Coo	CATION	n of the	

GENERATION OF PHOSPHORUS ESTER, PHOSPHORUS AMIDE, AND PHOSPHINE DERIVATIVES OF LOW COORDINATION NUMBER

FINAL REPORT

LOUIS D. QUIN

OCTOBER 15, 1986

U. S. Army Research Office
Post Office Box 12211
Research Triangle Park, NC 27709
Grant Number: DAAL03-86-K-0028

Duke University Durham, NC 27706

APPROVED FOR PUBLIC RELEASE;
DISTRIBUTION UNLIMITED

TABLE OF CONTENTS

	Page
I.	Purpose of the Research Program 1
II.	Major Accomplishments
	A. Synthesis of a Phosphaalkene Generator
	B. Synthesis of Metaphosphate Generators3
III.	Publications Resulting from this Grant4
IV.	Participating Scientific Personnel4

ANNANA MAKKANA SPECIOS SOCIOSOS MANANAS



Access	ion	For		7
HT1:	es es .	1	<u>/</u>	
2	٠,٠	•		
	100	. }	П	
		1.20.0		
-				
· · ·				
• • •	•	.27		
		1.5	lodes	
		10.3	jor [−]	

Al

I. Purpose of the Research Program

The 7-phosphanorbornene system is synthetically readily approached, and because of the considerable strain involved in the bridged structure, along with a mechanism provided by the double bond for a retro-cycloaddition process, this system was considered as a potential precursor of a variety of phosphorus species with low coordination numbers. The ring strain is known to permit the insertion of oxygen into a C-P bond, thus creating another ring system potentially capable of undergoing a retro[4+2] cycloaddition to release a phosphoryl compound of low coordination. A further extension of this concept could result in the synthesis of species with C-P double bonds, from the phosphabicyclo[2.2.2]octene system. To explore these possibilities, syntheses of suitable precursors are required. This report summarizes work both on the precursor syntheses, as well as on the successful generation of certain low-coordination species.

This is a report covering only the first seven months of a three-year grant period, and hence the accomplishments during this brief period are not numerous. This report is required by the transfer of the grant from Duke University to the University of Massachusetts.

II. Major Accomplishments

A. Synthesis of a Phosphaalkene Generator

15773432 (20100000

In previous work on this project, the synthesis of compounds $\underline{1}$ and $\underline{2}$ was accomplished. Preliminary work on their reduction to the corresponding phosphines $\underline{3}$ and $\underline{4}$ was not very successful, as decomposition of the phosphines accompanies their formation. We have now discovered conditions that allow the isolation of phosphine $\underline{3}$. The reduction may be smoothly

accomplished with trichlorosilane in chloroform at 15° or in benzene at 0° . These very mild conditions prevent the premature fragmentation of the phosphine, which ends up with the formation of Me₂PH and dimethyl phthalate. Phosphine $\frac{3}{2}$ has $\frac{5}{2}$ 31 P $^{-37}$ and gives the expected 13 C NMR spectrum. It is now available for studies on its thermal degradation, which is expected to proceed by a retro-cycloaddition to form CH₃P=CH₂ and the phthalate. The phosphaalkene will be highly reactive, and its formation must be confirmed by trapping experiments. Attempts to observe it in the gas phase at low pressure by photoelectron spectroscopy will be made in collaboration with Prof. N. Werstiuk, McMaster University.

Phosphine 4 has also been formed but in poor yield. It is much less stable and has not been fully characterized. It will be less useful as a precursor of a phosphaalkene.

$$\frac{0}{Ph-P}$$

$$\frac{HSiCl_3}{2}$$

$$\frac{4}{4}$$

B. Synthesis of Metaphosphate Generators

The sequence below is now well established as a means for generating the highly reactive metaphosphate species (5).

In continuing our studies on this process and on the properties of metaphosphates, we have carried out the synthesis of the three new compounds $\underline{6}$, $\underline{7}$, and $\underline{8}$. The first two should provide a precursor after the O-insertion reaction for metaphosphates that might be stabilized by the large bulky substituent on O. Compound $\underline{8}$ is of interest in providing a metaphosphate with a highly desirable O-substituent in the event that the metaphosphate finds utility as a phosphorylating agent in the nucleotide field.

The very larger tri-t-butylphenyl substituent of 6, however, completely prevented the O-insertion reaction, and no way has been found to overcome this problem. The synthesis of the mesityl derivative 7 has now been accomplished but the O-insertion has not yet been attempted. Compound 8 behaved well in the O-insertion process, although it proved to be surprisingly sensitive to water. Its decomposition to CCl₃CH₂OPO₂ proceeded smoothly and this species was detected by the usual device of trapping with alcohol or N-methylpyrrole.

III. Publications Resulting from this Grant

 L. D. Quin and G. Keglevich, "Stereochemistry of the Reaction of Oxygen Nucleophiles with a Phosphinous Chloride in the 7-Phosphanorbornene Series," J. Chem. Soc. Perkin Trans. II, 1029 (1986).

RECO TEXASES EXCERCE MAINTEE VANDANIA CONTRACTO ANNOVANA MAINTANA CONTRACTOR

2. L. D. Quin and J. Szewczyk, "7-Phosphanorbornene Derivatives as

Precursors of Esters and Amides of Two-coordinate Thiophosphenous O
Acid," J. Chem. Soc. Chem. Commun., 844 (1986).

IV. Participating Scientific Personnel

- 1. L. D. Quin, Research Director, 15 December 1985 31 July 1986
- Jerzy Szewczyk, Postdoctoral Research Associate, 1 January 1986 30
 June 1986.
- Bela Pete, Postdoctoral Research Associate, 1 April 1986 31 July
 1986.

IV. Participating Scientific Personnel (cont.)

TOTAL PROPERTY CONTROL PROPERTY SERVICES PROPERTY TOTAL PROPERTY DESCRIPTION OF THE PROPERTY O

- 4. Janell Boyd, Secretary, 20 December 1985 14 March 1986.
- 5. Robert Topping, Graduate Student, Research Assistant, 1 January 1986 31 March 1986
- 6. Brian Marsi, Graduate Student, Research Assistant, 1 May 86 31 August 1986
- 7. Xiao-Ping Wu, Graduate Student, Research Assistant, 1 May 1986 31 July 1986
- 8. Adrienne Lawler, Undergraduate Student, 12 May 1986 8 June 1986